

Fig 1. In vitro association of GEF-H1 with PAK4 alleles and subdomains.




Fig 2. PAK4 and GEF-H1 co-associate.




Fig 3. PAK4 phosphorylates GEF-H1 in vitro and in vivo.




Fig 4. Morphological effects of GEF-H1 alleles.




Fig 5a. PAK4 effects on the actin cytoskeleton

b




Fig 5b. PAK4 regulates lammellipodia formation through phosphorylation of S810.




Fig 6a. PAK4 signalling through GEF-H1M to the actin cytoskeleton.

b




Fig 6b. PAK4 activity destabilizes GEF-H1M association with MTs.




Fig 7. Model for reciprocal regulation of Rac and Rho *in vivo*.

Title: GEF-H1: BIOMARKERS,
COMPLEXES, ASSAYS AND
THERAPEUTIC USES THEREOF

Inventor(s): Tod R. SMEAL, et al.

Appl. No.: Unassigned

Phosphorylation scores GEF-H1 proteins and peptides.

FIGURE 8

a.

	PEPTIDE		PHOSPHORYLATION
681	RRKSLVGTPYWMAPE	PAK4 act loop	+++++
1008	RRRSLPAGDALYLSFNPP	GEFH1 (807-824)	+++++
1009	RRRSLPAGDALYLSFNPP	GEFH1 (807-824)	-
1010	RRRSLPAGDALYLSFNPP	GEFH1 (807-824)	+++++
1412	RQSLGSRGRS SLSLAK	GEFH1 (55-72)	-
1413	RQSLGSRGRS SLSLAK	GEFH1 (55-72)	-
1414	RQSLGSRGRS SLSLAK	GEFH1 (55-72)	+++++

b.

POLYPEPTIDE	PHOSPHORYLATION
1-386	++++
386-921	++
763-921	++++
51-921 ^{S810A}	++++
386-921 ^{S810A}	-
807-824	++++
55-72	++

Title: GEF-H1: BIOMARKERS,
COMPLEXES, ASSAYS AND
THERAPEUTIC USES THEREOF

Inventor(s): Tod R. SMEAL, et al.

Appl. No.: Unassigned

FIGURE 9

GEF-H1 regulatory regions are conserved with Cdc24

-----PFTQLAYCSSEVISE--ROSELLSOKQOEELKSNCAARD
SLRSKTTIRERPSSAIYPSSESFRSLLGSRRCRSSLSLAKSVSTTN

-PR-RXSLXG..

Consensus

Cdc24 (62-99)

GEF-H1M and S

QTH--DSMASFSSSHMKRVSDVLPK--RRRTSSSFSEIKSTSEN
GMEPLPNEAPWARRPVE-----PR-RRLPAG--DALYLSFNPPQPSRGTD

-PR-RXSLXG..

Consensus

Cdc24 (712-752)

GEF-H1M and S




Figure 10